

# The Benefits and Risks of Cannabis and Cannabinoids

Daniel J. Clauw M.D.

[dclauw@umich.edu](mailto:dclauw@umich.edu)

Professor of Anesthesiology, Medicine  
(Rheumatology), and Psychiatry  
Director, Chronic Pain and Fatigue  
Research Center

The University of Michigan

# Disclosures

- Consulting
  - Pfizer, Forest, Eli Lilly, Pierre Fabre, Cypress Biosciences, Wyeth, UCB, AstraZeneca, Merck, J & J, Nuvo, Jazz, Abbott, Cerephex, Iroko, Tonix, Theravance, IMC, Zynerba, Sammumed, Aptinyx
- Research support
  - Pfizer, Cypress Biosciences, Forest, Merck, Nuvo, Cerephex
- Testifying on behalf of State of Oklahoma against opioid manufacturers
- Went to the University of Michigan in the 1970's

# Benefits and Risks of Cannabinoids

- Definitions and Background
- Benefits of Cannabinoids
- Risks of Cannabinoids
- Role in Treating Chronic Pain
- Summary

# Benefits and Risks of Cannabinoids

- Definitions and Background
- Benefits of Cannabinoids
- Risks of Cannabinoids
- Role in Treating Chronic Pain
- Summary

# Definitions

- Cannabis – A genus of flowering plants with three different species: indica, sativa, and ruderalis
  - Can be bred to have low amounts of psychoactive compounds (e.g. THC) that are used to make hemp, or high amounts that are used for recreational/medicinal purposes
  - Sativex is a oral spray that is a cannabis extract
- Cannabinoid – Compounds that act at cannabinoid receptors
  - Endocannabinoids – endogenous ligands produced naturally that bind to CB1 and CB2 receptors
  - Phytocannabinoids – plant origin (cannabis/marijuana)
    - At least 80 different cannabinoids in cannabis
  - Synthetic cannabinoids

# Endocannabinoid system - I

**A set of receptors and their naturally occurring ligands and enzymes regulating control**

- **Receptors** – G-coupled protein receptors (the most abundant in CNS in man) on presynaptic membrane of cells in peripheral and central nervous system
  - CB1 – Primarily in central nervous system (but not in medulla in man) these act primarily to inhibit release of neurotransmitters
  - CB2 – Largely found in periphery on immune and nerve cells (although some in CNS on microglia and DRG)
  - Other receptors can bind these ligands because there is activity in CB1/CB2 knockouts (TRPV1, GPR55)

CB, cannabinoid receptor; CNS, central nervous system, DRG, dorsal root ganglion

Pertwee RG. Handb Exp Pharmacol. 2005;(168):1-51.


# Endocannabinoid system - II

- **Ligands** – Endocannabinoids are eicosanoid lipid messengers that are the physiological ligands for the cannabinoid receptors:
  - ananamide (N-arachidonylethanolamide, AEA)
  - 2-arachidonoylglycerol (2-AG)
  - PEA, virodamine, OAE
- **Enzymes** that synthesize and degrade the lipids endocannabinoids, such as fatty acid amide hydroxylase or monoacylglycerol lipase
  - Drugs being developed for pain that inhibit these enzymes

AEA, arachidonylethanolamine PEA, palmitoylethanolamide; OAE, oleoylethanolamine

Pertwee RG. Handb Exp Pharmacol. 2005;(168):1-51.

# Endocannabinoid system



Aizpurua-Olaizola, Oier, et al. "Targeting the endocannabinoid system: future therapeutic strategies." *Drug discovery today* 22.1 (2017): 105-110. Fonseca et al, *Prostaglandins & other lipid mediators* 102 (2013): 13-30.


# Endocannabinoid system - III

## Some known functions of the endocannabinoid system in humans:

- **Memory** – Generally affect short term memory, may play adaptive role in extinction of old memories in hippocampus
- **Neurogenesis**
- **Appetite** – Act in hypothalamus to increase appetite, inversely related to leptin levels
- **Analgesia**
- **Immune function** – Generally inhibit immune function, generally mediated via CB2<sup>1</sup> but some evidence CB1 might play role in T-cell responses. May be upregulation of CB2 receptors in some inflammatory disorders
- **Stress** – Help habituate/reduce HPA axis activity during repeated stress<sup>2</sup>

CB, cannabinoid receptor; HPA, hypothalamic-pituitary-adrenal

1. Rom S, Persidsky Y. J Neuroimmune Pharmacol. 2013; 8:608-20. 2. Hill MN, et al. Proc Natl Acad Sci U S A. 2010;107:9406-11.



CB, cannabinoid receptor  
 Image source: © Crowe S, et al. Brain Behav Immun  
 Crowe S, et al. Brain Behav Immun. 2014;42:1-5.

# Cannabis-derived cannabinoids

More than 80 known, with different strains having different relative concentrations

- **THC** (Synthetic forms include Dronabinol, Marinol, Nabilone)
  - The primary psychoactive cannabinoid in cannabis, and its metabolites are those assayed for in drug tests
  - Although it binds relatively equally to both the CB1 and CB2 receptors, most of its effects are associated with CB1 activity in brain
- **Cannabidiol (CBD)**
  - Is not psychoactive and does not bind to CB receptors, but has anticonvulsant and anti-inflammatory effects
  - May act as an indirect antagonist of CB agonists as thus has anti-psychotic effects
  - Also acts as 5HT1a agonist which might be responsible for potential analgesic, antidepressant effects

5HT1a, 5-hydroxytryptamine 1A receptor; CB, cannabinoid receptor; THC, tetrahydrocannabinol

Pertwee RG. Handb Exp Pharmacol. 2005;(168):1-51.

# Cannabis-derived cannabinoids

**More than 80 known, with different strains having different relative concentrations**

- **THC** (Synthetic forms include Dronabinol, Marinol, Nabilone)
  - The primary psychoactive cannabinoid in cannabis, and its metabolites are those assayed for in drug tests
  - Although it binds relatively equally to both the CB1 and CB2 receptors, most of its effects are associated with CB1 activity in brain
- **CBD** (cannabidiol)
  - Seems to be generally very well tolerated
  - Is not psychoactive and does not bind with any significant affinity to CB receptors, but yet has anticonvulsant and anti-inflammatory effects
  - Is actually thought to potentially protect against psychoactive effects of THC and hypothesized by some to be an effective anti-psychotic (although a recent Cochrane review concluded there was insufficient evidence of this)

# Benefits and Risks of Cannabinoids

- Definitions and Background
- Benefits of Cannabinoids
- Risks of Cannabinoids
- Role in Treating Chronic Pain
- Summary

# Potential Benefits of Cannabinoids

- Antiemetic<sup>1</sup> – Marinol is FDA-approved (Schedule III) for use in post-chemotherapy nausea/vomiting
- Anorexia – Marinol is FDA-approved for this use in AIDS-induced anorexia in US
- Anti-spasticity agent<sup>2</sup>
- Anticonvulsant<sup>3</sup> – Focus on CBD effects
- Neuroprotective
  - Being studied in Alzheimer's<sup>4</sup> because preclinical models show CB1/2 activation leads to reduction in beta-amyloid
  - Retrospective study of patients admitted with severe TBI showed significant reduction in death in those who had a positive drug screen for THC<sup>5</sup>
- Anti-tumor effects<sup>6</sup>

AIDS, acquired immune deficiency syndrome; CB, cannabinoid receptor; CBD, cannabidiol; FDA, Food and Drug Administration; TBI, traumatic brain injury; THC, tetrahydrocannabinol

1. Sharkey K, et al. Eur J Pharm. 2014; 722:134-46. 2. Koppel, et al. Neurology. 2014;82:1556-63. 3. Devinsky O. Epilepsia;2014;55:791-802. 4. Aso E, et. al. Front Pharmacol. 2014;5:37. 5. Nguyen BN. Am Surg. 2014;80:979-83. 6. Cridge B, Rosengren RJ. Cancer Manag Res. 2013;5:301-13.

# Anti-inflammatory effects

- Activation of CB2 receptors on immune cells leads to a variety of anti-inflammatory effects
- These (largely) peripheral effects can also lead to analgesic responses where pain in animal models or human diseases where pain is due to inflammation
- Ajulemic acid (Resunab) is a synthetic cannabinoid and the current formulation has greater selectivity for CB2>>CB1 and also activates PPAR-gamma. Thus it is considered relatively “non-psychoactive” and is in ongoing trials in autoimmune disorders and neuropathic pain

CB, cannabinoid receptor; PPAR, peroxisome proliferator-activated receptor

Boychuk DG, et al. J Oral Facial Pain Headache. 2015;29:7-14.

# Benefits and Risks of Cannabinoids

- Definitions and Background
- Benefits of Cannabinoids
- Risks of Cannabinoids
- Role in Treating Chronic Pain
- Summary

# Risks of Cannabinoids


- Almost all available data is from long term recreational users so we probably have good “worst case” data
- Partly related to route of administration
  - Smoking cannabis may lead to chronic bronchitis and potentially cancer of the mouth, throat, lung
  - This is likely reduced or eliminated with use of vaporizers or e-cigarettes
  - Oral administration causes less “likability” than inhalation or smoking and presumably no risk of bronchitis or cancer
  - *Individuals using cannabis for medicinal purposes should probably be using an oral formulation but dosing is problematic*
- The few deaths associated with cannabis are generally due to severe paranoia or tachycardia associated with overdose via oral administration

# Long Term Risks of Cannabinoids<sup>1</sup>

- Psychotic illnesses
  - It is now generally accepted that individuals who begin smoking cannabis prior to age 25 have 1.5 – 2.4X the rate of developing a psychotic illness<sup>2</sup>
  - This risk is modified by childhood trauma, family history of a psychotic illness, and perhaps genetic polymorphisms
- Long term effects on memory and brain structure
  - Both neuropsychological testing, and functional and structural neuroimaging studies, have suggested that individuals who use cannabis recreationally beginning in adolescence have decreased cognitive performance<sup>1,3</sup>
  - These studies have significant methodological issues because of other common exposures (e.g. alcohol or other illicit drugs) and behavioral issues in these individuals<sup>3</sup>

# Risks of Cannabinoids<sup>1</sup>

- Respiratory
- Dependence
  - Occurs in approximately 9% of individuals who use cannabis, but is about double in those who begin using in adolescence
  - This is lower than almost all other drugs of abuse (nicotine 32%, opioids 23%, alcohol 15%)
  - Highest risk in those with poor academic achievement, deviant behavior in childhood, poor parental relationships, family history of substance abuse
  - Physical addiction and withdrawal are much less common/severe than other drugs of abuse



### Comparison of classification systems for the harms and risks of drug abuse in the development of the multi-category Nutt rational scale

Correlation between mean scores from the independent experts and the specialist addiction psychiatrists 1=heroin. 2=cocaine. 3=alcohol. 4=barbiturates. 5=amphetamine. 6=methadone. 7=benzodiazepines. 8=solvents. 9=buprenorphine. 10=tobacco. 11=ecstasy. 12=cannabis. 13=LSD. 14=steroids

# Is there a link between marijuana and cancer?

- Smoked marijuana delivers THC and other cannabinoids to the body, but it also delivers harmful substances to users and those close by, including many of the same substances found in tobacco smoke, which are harmful to the lungs and cardiovascular system.
- Researchers have found limited evidence of an association between current, frequent, or chronic marijuana smoking and testicular cancer (non-seminoma-type)

# Cannabis and motor vehicle accidents

- Driving while impaired by any substance, including marijuana, is dangerous. Marijuana, like alcohol, negatively affects a number of skills required for safe driving:
  - Marijuana can slow your reaction time and ability to make decisions.
  - Marijuana use can impair coordination, distort perception, and lead to memory loss and difficulty in problem-solving.
- The risk of impaired driving associated with marijuana in combination with alcohol appears to be greater than that for either by itself.
- Latest statistics in states that have legalized cannabis suggests very small increase in MVA (3%) but no increase in fatalities

# Benefits and Risks of Cannabinoids

- Definitions and Background
- Benefits of Cannabinoids
- Risks of Cannabinoids
- Role in Treating Chronic Pain
- Summary

# Systematic reviews with meta-analysis on cannabis-based medicines for chronic pain: a methodological and political minefield

Winfried Häuser<sup>a</sup>, Nanna B. Finnerup<sup>b,c</sup>, R. Andrew Moore<sup>d</sup>

Public interest in cannabis products for medical purposes has been widely advocated, with legalization for recreational and medical use in North America and some European countries.<sup>6,8</sup> Legalization of cannabis-based medicines (CBMs) (medical cannabis, plant-based cannabinoids [tetrahydrocannabinol, cannabidiol, and combinations], and synthetic tetrahydrocannabinol analogues) has bypassed usual drug regulatory procedures.<sup>6</sup> Systematic reviews with meta-analyses of randomised controlled trials (RCTs) with CBM for chronic pain conditions help determine “post hoc” whether the preconditions of drug agencies for approval were met and to guide physicians and patients.

A systematic review of systematic reviews on CBM highlighted the uncertainty about whether CBMs improve pain, with only low or very low quality evidence available.<sup>1</sup> Individual systematic reviews generally avoided issues of trial quality, usually had some flaws, and included different drugs, doses, durations, conditions, and outcomes. Most reviews agreed that there was no, or no clinically relevant, effect.


In this issue of *PAIN*, Stockings et al.<sup>12</sup> provide the most comprehensive systematic review with meta-analysis of RCTs and observational studies with CBM, including 47 RCTs with 4271 patients with chronic noncancer pain. Study duration ranged between 1 day and 26 weeks. The authors avoid some methodology flaws of some of the systematic reviews mentioned above. They included “gray” literature and used all studies providing data in quantitative analyses. Average pain intensity, 30% and 50% or more pain relief, emotional and physical

There remains a methodological minefield, through which we need to step carefully. For example:

- (1) Most studies analysed are of low methodology quality.
- (2) Most studies included fewer than 50 patients per treatment arm. Small CBM studies are often the most positive.
- (3) Short-duration experimental studies (hours, a single day) were included, unhelpful in judging longer-term efficacy.
- (4) Lumping all chronic pain syndromes together does not help in managing individual patients, given the heterogeneity of chronic pain and its mechanisms. Even the importance of subgroup analyses is limited: cancer pain might have nociceptive and/or neuropathic components; neuropathic pain can have many dimensions, and drugs might be effective for some dimensions of neuropathic pain but not for others.<sup>3</sup> Whether heterogeneity of pain mechanisms is relevant for the efficacy of CBM, which are nonspecific centrally acting drugs is, however, unknown.
- (5) Lumping together all CBMs, including experimental drugs unavailable for clinical use, limits the clinical relevance of combined results.
- (6) There is the risk of overestimating the effects of CBM for pain relief because of unpublished studies, for example, with nabilone for chronic neuropathic pain.<sup>10</sup>
- (7) Long-term risk and severe but rare side effects are not captured in small, short-duration trials.

What can patients, clinicians, trialists, drug companies, and

# Cannabis clinical trials for chronic pain



- Limited: short length and small sample size
  - Many used THC alone or THC + CBD
- Most support for use of cannabinoids in neuropathic pain (THC+CBD).
- Increased risk of short term AEs (mostly minor) for study participants


Whiting, Penny F., et al. *Jama* 313.24 (2015): 2456-2473.

Nugent, Shannon M., et al. *Annals of internal medicine* 167.5 (2017): 319-331.


# Anti-inflammatory effects of CBD

- There are many animal models where CBD has been demonstrated to have potent anti-inflammatory effects in a variety of models (including murine collagen arthritis and carrageenan models) but it is much less clear how those anti-inflammatory effects are being mediated
- Some evidence that anti-inflammatory effects might be occurring via CB2 (very high doses needed), adenosine receptors, arachidonic acid release (causes shift from cyclooxygenase to lipoxygenase pathway), via direct inhibition of cytokine production, or via binding to the GPR55 receptor (which has both inflammatory and nociceptive properties)

# ZYN002 - Median Weekly Average Worst Knee Pain Score over Time - Males



## ZYN002 - Median Weekly Average Worst Knee Pain Score over Time – Females



# Mechanistic Characterization of Pain

## Variable degrees of any mechanism can contribute in any disease

	Nociceptive	Neuropathic	Centralized
<b>Cause</b>	Inflammation or damage	Nerve damage or entrapment	CNS or systemic problem
<b>Clinical features</b>	Pain is well localized, consistent effect of activity on pain	Follows distribution of peripheral nerves (i.e. dermatome or stocking/glove), episodic, lancinating, numbness, tingling	Pain is widespread and accompanied by fatigue, sleep, memory and/or mood difficulties as well as history of previous pain elsewhere in body
<b>Screening tools</b>		PainDETECT	Body map or FM Survey
<b>Treatment</b>	NSAIDs, injections, surgery, ? opioids	Local treatments aimed at nerve (surgery, injections, topical) or CNS-acting drugs	CNS-acting drugs, non-pharmacological therapies
<b>Classic examples</b>	Osteoarthritis Autoimmune disorders Cancer pain	Diabetic painful neuropathy Post-herpetic neuralgia Sciatica, carpal tunnel syndrome	Fibromyalgia Functional GI disorders Temporomandibular disorder Tension headache Interstitial cystitis, bladder pain syndrome




CNS, central nervous system; FM, fibromyalgia; GI, gastrointestinal

Clauw DJ. The taxonomy of chronic pain: Moving towards more mechanistic classification. In: Wallace DJ & Clauw DJ, editors. Fibromyalgia and other central pain syndromes. Philadelphia: Lippincott, Williams & Wilkins; 2005. p.10-16.

# Neural Influences on Pain and Sensory Processing

## Facilitation

- Substance P ↑
- Glutamate and EAA ↑
- Serotonin (5HT<sub>2a, 3a</sub>) ↑
- Nerve growth factor ↑



## Inhibition

- Descending anti-nociceptive pathways ↓
- Norepinephrine-serotonin (5HT<sub>1a,b</sub>), dopamine ↓
- Opioids ↑
- GABA ↓
- Cannabinoids ↓

5HT, 5-hydroxytryptamine/serotonin; EAA, excitatory amino acid; GABA, gammahydroxybutyrate  
1. Schmidt-Wilcke T, Clauw DJ. Nat Rev Rheumatol. 2011;7:518-27  
2. Clauw DJ. JAMA. 2014;311:1547-55.

# Centralization Continuum

*Proportion of individuals in chronic pain states that have centralized their pain*

Peripheral

Centralized



Acute pain

Osteoarthritis

SC disease

Fibromyalgia

RA

Ehler's Danlos

Tension HA

Low back pain

TMJD IBS


HA, headache; IBS, irritable bowel syndrome; SC, sickle cell disease; TMJD, temporomandibular joint and muscle disorders

Image source: © Clauw DJ. Philadelphia: Lippincott, Williams & Wilkins; 2005. p.10-16

Adapted from: Clauw DJ. The taxonomy of chronic pain: Moving towards more mechanistic classification. In: Wallace DJ & Clauw DJ, editors. Fibromyalgia and other central pain syndromes. Philadelphia: Lippincott, Williams & Wilkins; 2005. p.10-16.


# Pharmacokinetics

## Pharmacokinetics




Route of administration influences THC pharmacokinetics, left = 5 mg i.v. injection, center = smoking 13.0 mg, or right = consuming cookie with 20 mg (Agurell et al. 1986).

# Feelings of 'high' from different administration routes



# U-Shaped Curve for cannabis effects



1. Hill KP. *Jama*. 2015;313(24):2474. 2. Wallace M, Schulteis G, Atkinson JH, Wolfson T, Lazzaretto D, Bentley H, et al. *Anesthesiology*. 2007;107(5):785–96. 3. Portenoy RK, Ganae-Motan ED, Allende S, Yanagihara R, Shaiova L, Weinstein S, et al. *J Pain*. Elsevier Ltd; 2012;13(5):438–49.


# Treating Based on Mechanisms

*Any combination may be present*

	Peripheral (nociceptive)	Neuropathic	Centralized Pain
NSAIDs	+	-	-
Opioids	+	+	-
Surgery/ Injections	+	+	-
Tricyclics	+	+	+
SNRIs	+	+	+
Gabapentinoid	-	+	+
CBD	+	-	-
THC	-	+	+

# Cannabis as an opioid substitute for chronic pain?

- Cannabis as a synergist with opioids?<sup>1,2</sup>
- State-wide analyses<sup>3-5</sup>
  - Importance of Dispensaries in these studies (Powell et al, 2018)
- Cross-sectional<sup>6-8</sup> and longitudinal support<sup>9-11</sup>




1. Elikottil, Jaseena, et al. *Journal of opioid management* (2009). 2. Abrams et al, *Clinical Pharmacology and Therapeutics*, (2011). 3. Bachhuber MA et. al. *JAMA Int Med* (2014). 4. Bradford and Bradford *Health Affairs*, (2016) 5. Bradford and Bradford, *Health Affairs* (2017). 6. Boehnke, Kevin F., Evangelos Litinas, and Daniel J. Clauw. *The Journal of Pain* (2016). 7. Lucas et al, *Journal of International Drug Policy* (2017) 8. Reiman et al, *Cannabis and Cannabinoid Research* (2017). 9. Haroutounian et al., *Clinical Journal of Pain* (2016). 10. Stith et al, *PLOSone* (2017) 11. Abuhasira et al, *European Journal of Internal Medicine*, (2018)

# Proposed marketing program for medical cannabis

Cannabis plant talking to opium producing poppy plant



We don't  
suck as  
bad as you  
do



# Benefits and Risks of Cannabinoids

- Definitions and Background
- Benefits of Cannabinoids
- Risks of Cannabinoids
- Role in Treating Chronic Pain
- Summary

# Pragmatic Advice for Using Cannabinoids in 2019

- Where possible use a cannabinoid or cannabinoid extract of consistent and known potency
- Start with CBD alone and then go to low dose of low THC:high CBD strain and go up slowly
- Emerging evidence of U-shaped curve
- Oral dosing better once stable dose and strain identified
- The strongest recommendation based on current benefit: risk data is for the use of cannabinoids instead of opioids for neuropathic or centralized pain states
  - Data from US suggest that legalizing cannabis in a state leads to fairly dramatic reductions in opioid overdoses<sup>1</sup>
- Use with caution in individuals under age 25

CBD, cannabidiol; THC, tetrahydrocannabinol

1. Bachhuber MA, et. al. JAMA Int Med 2014;174:1668-73.

# Summary

- The endocannabinoid system is widely distributed in the human body and there is strong biological plausibility that these can be effective and safe analgesics at the right dose and in the right person
- Cannabinoids can exert analgesic effects in the periphery (mainly anti-inflammatory) and CNS (dissociate individuals from the sensory experience of pain)
- For treatment of clinical pain with cannabinoids there is best evidence that they are effective in neuropathic and centralized pain, and little current evidence that they are effective in nociceptive/inflammatory pain states

# Pragmatic Advice for Using Cannabinoids in 2019

- Where possible use a cannabinoid or cannabinoid extract of consistent and known potency
- Start with CBD alone and then go to low dose of low THC:high CBD strain and go up slowly
- Emerging evidence of U-shaped curve
- Oral dosing better once stable dose and strain identified
- The strongest recommendation based on current benefit: risk data is for the use of cannabinoids instead of opioids for neuropathic or centralized pain states
  - Data from US suggest that legalizing cannabis in a state leads to fairly dramatic reductions in opioid overdoses<sup>1</sup>
- Use with caution in individuals under age 25

CBD, cannabidiol; THC, tetrahydrocannabinol

1. Bachhuber MA, et. al. JAMA Int Med 2014;174:1668-73.

# Cannabis and Cannabinoids

- Definitions and Background
- Overview of Risks and Benefits of Cannabinoids
- Role in Pain Management
  - Acute vs. Chronic
- Summary

# What to do when candidate for surgery discloses cannabis use?

- No need to alter typical analgesic regimen.
- Use best judgment
- No evidence for cannabinoids in surgical setting or for acute pain.
- Given length of in-patient stay, don't really do anything for any user if patients are there for more than 1-2 days.

# Summary

- Endocannabinoid system: plausibility as analgesic at right dose and in right person
- Cannabinoids can exert analgesic effects in the periphery and CNS
- For treatment of clinical pain with cannabinoids:
  - Better evidence for efficacy in chronic neuropathic pain: May be reasonable to use in place of opioids for *chronic pain* when all else has failed, but NOT in a surgical context.
  - No evidence for efficacy in acute/surgical pain. No clinical trial data on recovery from surgical pain

**Questions?**